

Liovital® *mirtillo*

8 SINGLE-DOSE MICROVIALS
FOR ORAL USE WITH RESERVOIR CAP



Food supplement

Vaccinium myrtillus L. dry extract

Vitis vinifera L. dry extract

Micronized pollen

Freeze-dried royal jelly

GLUTEN FREE

INGREDIENTS

Content of the reservoir cap: freeze-dried royal jelly; maltodextrin; anticaking agents: silicon dioxide and magnesium oxide; bulking agent: microcrystalline cellulose.

Content of the microvial: water; fructose; dehydrated concentrate blueberry (*Vaccinium myrtillus L.*) juice (blueberry, maltodextrin); blueberry (*Vaccinium myrtillus L.*) fresh berries dry extract std. 1% in anthocyanin expressed as cyanidin 3-glucoside chloride (blueberry, dextrose); red grapevines (*Vitis vinifera L.*) dry seeds extract std. 95% in procyanidin; water-soluble micronized pollen; acidity regulator: sodium lactate solution 60%; preservative: potassium sorbate; mixed berries flavour, raspberry flavour; acid: citric acid.

PACKAGING AND CONTENTS

Box containing 8 single-dose microvials in amber PET, of 10 ml each, with reservoir cap.

PRODUCT CATEGORY

Food supplement.

INDICATIONS

LIOVITAL® *mirtillo* contains standardized and titrated extracts of blueberry (fruits) and red grapevines (seeds) associated with water-soluble micronized pollen and with freeze-dried royal jelly.

Blueberry is useful for microcirculation efficiency; pollen and royal jelly are rich in essential amino acids, vitamins and minerals.

RECOMMENDED DOSE

1 or 2 microvials per day, based on needs.

STORAGE CONDITIONS

Store at room temperature (15 – 30°C) and do not expose to heat sources. The expiry date refers to the product unopened and correctly stored.

WARNINGS

Do not exceed the recommended daily dose. Food supplements are not intended as a substitute of a balanced and varied diet and a healthy lifestyle. Keep out of reach of children under three years old.

NOTICE: since royal jelly is a colloidal and not-soluble substance, the reconstituted microvial is presented as a lactescent suspension.

INSTRUCTIONS FOR USE

- 1) Screw completely the cap until the breaking of the seal, so the powder in the capsule will fall into the microvial.
- 2) Shake well before use.
- 3) Unscrew the cap.
- 4) Take orally according to instructions.

The persistence, after shaking step, of undissolved particles is an integral part of the product characteristics and should not be understood as a sign of alteration.



PRODUCED AND PACKAGED:

see external packaging.

INTERNAL LEAFLET LIOVITAL MIRTILLO MICROVIALS - REV.8 OF 12/2019

Pharma Line S.r.l.- Via Bertani 2 - 20154 Milan, Italy.

www.pharma-line.it

R.0-0318